

Research Article

Effectiveness of an Exercise Protocol for Dynamic Balance in Amateur Female Soccer Players: A Randomized Clinical Trial.

Shirley Mireya Ortiz Pérez¹, David Marcelo Guevara Hernández^{1*}, Vanessa de los Ángeles Belloso de Noriega², Fernando Daniel Álvarez Maigualema³

¹ Departamento de Fisioterapia, Facultad de Ciencias de la Salud, Universidad Nacional de Chimborazo. Riobamba, Ecuador.

² Departamento de Enfermería, Facultad de Medicina, Universidad del Zulia. Maracaibo, Venezuela.

³ Departamento de Fisioterapia, Facultad de Enfermería, Pontificia Universidad Católica del Ecuador. Quito, Ecuador.

***Corresponding author:** David Marcelo Guevara Hernández,
Departamento de Fisioterapia, Facultad de Ciencias de la Salud, Universidad Nacional de Chimborazo. Riobamba, Ecuador.
E-mail: davidzguevara@icloud.com

ORCID ID:

Shirley Mireya Ortiz Pérez: 0000-0003-0648-2381

David Marcelo Guevara Hernández: 0000-0001-5063-0519

Vanessa de los Ángeles Belloso de Noriega: 0009-0000-5030-5409

Fernando Daniel Álvarez Maigualema: 0009-0002-8430-2657

Article info:

Received: 28 Augt 2024

Accepted: 18 Jan 2025

Running Title : Dynamic balance in amateur soccer players

Abstract

Background: Women's soccer has gained popularity in the last decade as well as preventive training strategies, therefore the present research aims to determine the effectiveness of a dynamic balance exercise protocol in amateur soccer players.

Materials and Methods: Twenty-four females participated in this study and were randomly assigned to a control group (12) and an experimental group (12). The experimental group received a dynamic balance exercise intervention with three sessions per week for six weeks, while the control group maintained their regular training. Balance was assessed before and after by the Y Balance Test in its anterior, posterior medial and posterior lateral directions. Data were analyzed through the T-test for independent and related samples.

Results: The analysis of this study shows significant improvements on dynamic balance $p=0.000$ in the experimental group after the intervention of the dynamic balance exercise program for six weeks, while in the control group there were no significant improvements.

Conclusions: Dynamic balance exercise program can be used to increase the balance of amateur female soccer players.

Keywords: Proprioception, Postural balance, Soccer, Exercise therapy

Introductions

Women's football is a discipline that demands a unique combination of physical and mental skills; speed, coordination, endurance, and agility are just some of the attributes players develop (1,2). However, beyond the sporting aspect, this discipline has proven to be a powerful driver of social change, promoting gender equality and empowering women worldwide. As a contact sport, it entails injuries associated with factors such as age, sex, player level, and prior injuries. It has been reported that women are at higher risk of sustaining knee and ankle injuries (3,4).

Regarding injuries and the player's level of experience, the risk appears to be heightened among younger players who practice the sport recreationally (5). Unlike professional athletes, amateur players balance their academic or work responsibilities with their football practice. This dual role results in less exclusive dedication to training, which translates into less specialized physical preparation and limited access to high-quality resources such as sports equipment and technical monitoring (6,7)

Injury prevention in football is essential for optimizing athletic performance (8). An injury-free player can train consistently and with greater intensity, leading to continuous improvement in technical and physical skills. Furthermore, avoiding long recovery periods minimizes the loss of physical condition and ensures greater availability for matches. The confidence derived from feeling safe and protected from injuries motivates players to perform at their best, which, in turn, benefits the team (9,10).

The importance of injury prevention programs for football players has been highlighted in previous research, emphasizing strategies such as balance training to improve proprioception, muscle strengthening, core-focused exercises, and specific variations of plyometric drills (11–13). Balance training through exercises involving diverse positions can modify afferent input and enhance proprioception, yielding positive effects on athletes' equilibrium. Previous studies (14,15) have demonstrated that this type of training improves balance in basketball players. However, there is limited evidence regarding such training in female football players. Therefore, the objective of this study was to determine the effectiveness of an exercise protocol for dynamic balance in amateur female football players (16).

Material and methods

Study design.

The design of this research is a prospective cohort study comprising a sub analysis of a randomized clinical trial to determine the effectiveness of the exercise protocol for dynamic balance of female soccer players. Through measurements taken before and after the intervention to 24 athletes aged 20 to 23 years, who met the inclusion requirements to subsequently form part of the experimental group (EG) and the control group (CG) whose allocation was randomized to eliminate bias. This research was executed for 6 weeks from May to June 2024. In compliance with the declaration of Helsinki.

Sampling method

The sampling was intensive to recruit eligible athletes. The population of this research beginner women footballers who have joined the university football team within the last 8 months and actively participate in training sessions of the soccer team at the Universidad Nacional de Chimborazo between the ages of 20 and 23 years old. A total of 40 soccer players were evaluated by means of a self-informed questionnaire. After the selection process, 24 players who fulfilled all the inclusion criteria were recruited for this research.

Study population and recruitment.

The inclusion criteria were: (1) female soccer players from female teams of the Universidad Nacional de Chimborazo, (2) athletes between 18 and 23 years old, (3) subjects free of injuries, (4) previous experience of at least one year practicing soccer, (5) with a training frequency twice a week (17). The following were excluded: (1) soccer players with a history of injury in the last six months, (2) with neurological or vestibular dysfunctions affecting balance control, (3) athletes with alterations in any of the joints of the lower extremities (15).

The research was socialized in person with the soccer players, and they were also informed of the study through their e-mails and cell phones. All the athletes were freely invited to be part of the research. Those interested received written information and it was emphasized that participation before and during the study was voluntary; each player could refuse or withdraw at any time during the study. Finally, before filling out the participation form, the volunteers had to read and understand all the information about the study, complete the form and register their consent to participate.

Investigation procedure

Twenty-four soccer players comprising the entire sample were randomly assigned into two groups, EG and CG. Both groups were intervened simultaneously, the average duration of each session was 45 minutes. The subjects of the experimental group performed a protocol of balance exercises with a frequency of three sessions per week, for six weeks, totaling eighteen sessions.

The control group, in contrast, participated in a training program comprising warm-up exercises, strength training, agility drills, sports-specific movements, and stretching exercises, maintaining the same duration, number of sessions, and frequency as the experimental group.

Prior to the intervention, all participants were verified for informed consent and resocialized about each study procedure. Furthermore, they were informed that they could voluntarily withdraw at any time from the research without giving any reason. To obtain the study results of each group, the athletes underwent the Y Balance Test, which was initially performed as a test with the aim of familiarizing each subject with the test to avoid errors that could affect the results. The test was performed at the beginning and at the end of the study.

Exercise protocol for balance.

The EG underwent the protocol of exercises for balance that was executed during their routine training, this protocol has a duration between 25 and 30 minutes, adding to the athlete's training a proprioceptive training to improve motor coordination in the central nervous system. The protocol is based on the postural change of the athlete by executing various positions as described by Lee et al, a) tandem, b) split squat, c) single leg, d) bilateral jump and landing, e) single leg jump and landing. All exercises increased in difficulty around the time of execution, the degree of effort of each athlete and the combination of exercises (*Figure 1*). The protocol was executed gradually in

each athlete to increase the proprioceptive effort, a determining factor to obtain a better balance through sensorimotor integration.




Figure 1. Balance Exercises

Research measurement instrument.

The balance of female soccer players was determined by the Y Balance Test (15), the athletes had to balance on only one of their lower limbs, while the limb in the air had to try to reach the farthest distance in several directions; a) anterior, b) posterolateral and c) posteromedial (*Figure 2*). Before taking the official measurements, the subjects performed a pretest.



Figure 2. Y Balance Test

Data collection.

The subjective and objective data of each athlete were collected through the clinical history. As for the dynamic balance was determined by the Y Balance Test these data were initially recorded in the Excel program from Microsoft 365 Office package, for subsequent analysis in the statistical package for social sciences "SPSS" version 29.

Statistical analysis

The measures obtained by the Y Balance Test were analyzed in SPSS through the T-test for independent samples to compare the difference in the dependent variable between the experimental group and the control group; and the T-test for related samples to compare the difference before and after the intervention in the experimental group with the objective of determining the effectiveness of the exercise protocol for the dynamic balance of beginner female soccer players. The significance level was predetermined at $p < 0.05$.

Results

The demographic data of the participants in this research are characterized by similarity in age, height, and weight, confirming a homogeneous group. The descriptive characteristics of the twenty-four participants are shown in *Table 1*.

Table 1. Demographic data of the experimental and control groups.

Variables	Experimental (n=12)	Group Control Group (n=12)
Age (years)	21.33 ±0.88	21.42 ±0.69
Height (cm)	155.92 ±4.64	154.08 ±4.73

Weight (kg)	59 ±9.06	57 ±4.30
--------------------	----------	----------

Table 2 shows the result of the comparison between the control group and the experimental group for dynamic balance in absolute reach before the intervention. The pre-intervention absolute values of dynamic balance in the control group (63.83 ± 8.08 cm) and the experimental group (75.29 ± 6.43 cm) revealed that there was no significant difference, showing that the group was homogeneous before the intervention ($p=0.002$). The intervention of the experimental group lasted 18 motor training sessions achieving a significant difference with an absolute range ($p=0.000^*$) between the control group (64.01 ± 7.85 cm) and the experimental group (81.76 ± 5.39 cm).

Table 2. Pre- and post-test of dynamic balance in the control and experimental groups in absolute range.

	Control Group	Experimental Group	P
Pretest	63.83± 8.08	75.29 ±6.43	0.002
Post test	64.01± 7.85	81.76 ±5.39	0.000*

The absolute results of the experimental group before 75.29 cm and after the intervention 81.76 cm during 18 motor training sessions show an increase of 6.47 cm. The results of the intervention in the experimental group show a significant improvement in dynamic balance in absolute reach ($p=0.000^*$) Table 3.

Table 3. Pre and post-test of dynamic balance in the experimental group in absolute range.

	Pre test	Post test	p
Experimental Group	75.29 ±6.43	81.76 ±5.39	0.000*

Discussion

Authors such as Benis et al. performed an intervention using body weight neuromuscular exercises for 8 weeks to improve postural control and stability of the lower extremities in female basketball players, with no history of injury. Where neuromuscular warm-up exercises were performed in the experimental group and in the control group a standard warm-up, the workouts were performed for 8 weeks, the exercises progressed from a stable to an unstable position to increase lower extremity strength demand and stability, while the control group performed aerobics, the authors added body weight plyometric exercises to the experimental group with the purpose of training the core muscles and activating the nervous system concluding that the scores on the Y Balance Test Postero medial improve in the experimental group, the same author alludes that the program also increases joint awareness and improves postural control (18).

On trunk stabilization exercises Filipa et al.; Granacher et al.; Imai et al.; D'souza et al. share programs based on core training or trunk stabilization through different sets of exercises (19–22). Filipa et al. describes as an objective to determine whether a neuromuscular training program focused on core stability and strength in the lower extremities would affect the performance of the balance test Star exclusion balance test (SEBT). The experiment was performed in 20 non-injured

soccer players by applying an 8-week program, due to the intervention in 16 sessions, there was evidence of improvement in the balance score measured with the SEBT especially in the posterolateral and posteromedial reach, this due to better neuromuscular control in the lower extremities (19).

Granacher et al. runs a 6-week intervention program comparing core strength training on unstable versus stable surfaces, achieving improvements on balance, coordination and speed, attributing its cause to the improvement on anticipatory muscle activation which is the basis for functional movements providing stability from the trunk to the lower extremities (20).

Imai et al. similarly performed trunk stabilization exercise and conventional trunk exercises to improve balance and strength, showing that trunk stabilization exercises improve static balance immediately and also sports performance. Adjudicating these improvements in the postero-lateral and posteromedial directions of the SEBT (21).

The primary finding of our study was that a 6-week balance training program, utilizing progressively challenging exercises, proved beneficial by yielding significant improvements in Y Balance Test measurements. This suggests that the specific exercise-based intervention had a positive effect on participants' balance compared to the control group, particularly regarding dynamic balance training. Therefore, the literature supports the use of trunk musculature training to improve balance, thus it is often found in a multitude of trainings to prevent ankle or knee injuries because decreased neuromuscular control of the trunk seems to influence the dynamic stability of the lower extremity during high-speed athletic maneuvers (23).

McKay et al. who conducted a treatment to evaluate the relationship between hip strengthening and pain in female runners with iliotibial band syndrome, through assigned groups which used conventional exercise, stretching, or strengthening of the hip muscles, the hypothesis was based on adding progressive complexity to the basic exercises for 8 weeks, the Y Balance Test was used to measure the differences in the patients, resulting that the stretching and hip training group improved the balance test mentioned in the patients, therefore it is also important to emphasize that functional exercise including sport specific movements improves balance and mobility (24).

Lee et al. conducted a six-week study, which, unlike ours, performed perturbations during the execution of the exercises and showed significance when measuring dynamic balance. The author indicates that proprioception is an important factor in maintaining dynamic balance and sensation of the limbs in space, especially necessary in basketball athletes who need dynamic balance that includes perturbation due to the nature of the game (25).

Our study was developed in female soccer players, a sport that requires constant movement changes and joint stability, so balance training is key to obtain mechanical advantages and minimize the overload or injury complexes on the ankle, although it is true that for now this study has focused on improving balance. In the future it would be interesting to project ourselves to the prevention of frequent injuries in women's soccer teams and detect possible predictions of lower body injuries based on balance.

Conclusions

The dynamic balance exercise program executed for six weeks with eighteen sessions promotes a significant improvement of balance in amateur female soccer players. This could contribute to injury prevention protocols; however, it is important to emphasize that more studies are needed that relate balance to the prevention of lower extremity injuries in women.

Study limitations

The study was limited by a small sample size.

Acknowledgments

The authors thank the owners of Neurofit Ec, for approval for the use of their laboratory and their support during the development of the manuscript.

Author's contributions

Conceptualization and supervision: Shirley Mireya Ortiz Pérez and David Marcelo Guevara Hernández, Methodology: Shirley Mireya Ortiz Pérez and David Marcelo Guevara Hernández, Data collection: Shirley Mireya Ortiz Pérez and Fernando Daniel Álvarez Maigualema, Data analysis: Vanessa de los Ángeles Belloso de Noriega and David Marcelo Guevara Hernández, Investigation and writing: Shirley Mireya Ortiz Pérez and David Marcelo Guevara Hernández.

Ethical Considerations

Compliance with ethical guidelines

This research was registered in the clinical trial database of the Neurofit ec physical rehabilitation center (CEINF00010) and was approved by the research ethics committee of the comprehensive physical rehabilitation center (N°2/2024 approval date February 11, 2024).

Funding

The authors did not receive financing for the development of this research.

Conflict of interest

The authors declare that there is no conflict of interest.

References

1. Gurau TV, Gurau G, Musat CL, Voinescu DC, Anghel L, Onose G, et al. Epidemiology of Injuries in Professional and Amateur Football Men (Part II). *J Clin Med.*2023;12(19):6293.
2. Moriyama M, Kouzaki M, Hagio S. Anticipatory postural control in adaptation of goal-directed lower extremity movements. *Sci Rep.* 2024;14(1):4142.
3. Crossley KM, Patterson BE, Culvenor AG, Bruder AM, Mosler AB, Mentiplay BF. Making football safer for women: a systematic review and meta-analysis of injury prevention programmes in 11 773 female football (soccer) players. *Br J Sports Med.* 2020;54(18):1089-98.
4. Kim MK, Shin YJ. Immediate Effects of Ankle Balance Taping with Kinesiology Tape for Amateur Soccer Players with Lateral Ankle Sprain: A Randomized Cross-Over Design. *Med Sci Monit Int Med J Exp Clin Res.* 2017;23:5534.
5. Alahmad TA, Kearney P, Cahalan R. Injury in elite women's soccer: a systematic review. *Phys Sportsmed.* 2020;48(3):259-65.
6. Bakarman MA, Tashkandi M, Mohammed NS. An intervention plan for preventing and handling amateur soccer players' injuries. *Niger J Clin Pract.* 2022;25(11):1816.
7. Pajonková F, Lenková R, Sučka J, Mikuláková WB. The impact of neuromuscular warm-up on the level of dynamic stability of lower limbs in female football players in the context of injury prevention. *Fizjoterapia Pol.* 2024;24(3):89-94.

8. Ramírez MAC, Oñate GH, Salamanca DEL. Características del pie y equilibrio dinámico en basquetbolistas, futbolistas y voleibolistas colombianos. *Rev Científica Salud Uninorte*. 29 de febrero de 2024;40(1):143-59.
9. Grygorowicz M, Wiernicka M, Wiernicka M. Systematic Review of Cost-Effectiveness of Injury Prevention Interventions in Soccer-Evidence Why Health Agencies Should Address It. *Int J Environ Res Public Health*. 2021;18(22):11901.
10. Pérez-Gómez J, Adsuar JC, Alcaraz PE, Carlos-Vivas J. Physical exercises for preventing injuries among adult male football players: A systematic review. *J Sport Health Sci*. 2022;11(1):115-22.
11. Ke W, Te H. Meta-analysis of meta-analyses of anterior cruciate ligament injury reduction training programs. *J Orthop Res Off Publ Orthop Res Soc*. 2018;36(10). Disponible en: <https://pubmed.ncbi.nlm.nih.gov/29737024/>
12. Syafrianto D, Randi TG, Alimuddin, Liza. The impact of core stability training on the dynamic balance of Solok city-based Saiyo FC soccer players. *J Phys Educ Sport*. 2023;23(12):3356-63.
13. Cè E, Longo S, Paleari E, Riboli A, Limonta E, Rampichini S, et al. Evidence of balance training-induced improvement in soccer-specific skills in U11 soccer players. *Scand J Med Sci Sports*. 2018;28(11):2443-56.
14. Sonesson S, Lindblom H, Hägglund M. Higher age and present injury at the start of the season are risk factors for in-season injury in amateur male and female football players—a prospective cohort study. *Knee Surg Sports Traumatol Arthrosc*. 2023;31(10):4618-30.
15. Lee AC, Sankaravel M, Chen ZX. The Effectiveness of Six-Week Balance Training with Perturbation Intervention in Enhancing Dynamic Balance of Basketball Players. *Phys Educ Theory Methodol*. 2023;23(1):21-7.
16. Al Attar WSA, Khaledi EH, Bakhsh JM, Faude O, Ghulam H, Sanders RH. Injury prevention programs that include balance training exercises reduce ankle injury rates among soccer players: a systematic review. *J Physiother*. 2022;68(3):165-73.
17. Lee UY, Joo CH, Lee UY, Joo CH. The effects of proprioceptive exercise training on physical fitness and performance of soccer skills in young soccer players. *J Exerc Rehabil*. 2024;20(1):34-41.
18. Benis R, Bonato M, Torre AL. Elite Female Basketball Players' Body-Weight Neuromuscular Training and Performance on the Y-Balance Test. *J Athl Train*. 2016;51(9):688-95.
19. FILIPA A, BYRNES R, PATERNO MV, MYER GD, HEWETT TE. Neuromuscular Training Improves Performance on the Star Excursion Balance Test in Young Female Athletes. *J Orthop Sports Phys Ther*. 2010;40(9):551-8.
20. Granacher U, Schellbach J, Klein K, Prieske O, Baeyens JP, Muehlbauer T. Effects of core strength training using stable versus unstable surfaces on physical fitness in adolescents: a randomized controlled trial. *BMC Sports Sci Med Rehabil*. 2014;6:40.
21. Imai A, Kaneoka K, Okubo Y, Shiraki H. EFFECTS OF TWO TYPES OF TRUNK EXERCISES ON BALANCE AND ATHLETIC PERFORMANCE IN YOUTH SOCCER PLAYERS. *Int J Sports Phys Ther*. 2014;9(1):47-57.
22. D'souza CJ, Santhakumar H, Bhandary B, Rokaya A. Immediate effect of stabilization exercises versus conventional exercises of the trunk on dynamic balance among

trained soccer players. Hong Kong Physiother J Off Publ Hong Kong Physiother Assoc Ltd Wu Li Chih Liao. 2022;42(1):23-30.

23. Imai A, Imai T, Iizuka S, Kaneoka K. A Trunk Stabilization Exercise Warm-up May Reduce Ankle Injuries in Junior Soccer Players. *Int J Sports Med.* 2018;39(4):270-4.
24. McKay J, Maffulli N, Aicale R, Taunton J. Iliotibial band syndrome rehabilitation in female runners: a pilot randomized study. *J Orthop Surg.* 24. 2020;15:188.
25. Lee HM, Oh S, Kwon JW. Effect of Plyometric versus Ankle Stability Exercises on Lower Limb Biomechanics in Taekwondo Demonstration Athletes with Functional Ankle Instability. *Int J Environ Res Public Health.* 2020;17(10):3665.