Research Article



Effects of Hamstring Flossing on Balance and Foot Function in Female Patients with Plantar Fasciopathy: A Randomized Controlled Study

Eman Ahmed Ahmed * 📵, Nagui Sobhi Nassif 📵, Rafik Elmaamoon Radwan 📵, Amal Abdelrahman Elborady 📵, Yassmin Essam Mohamed 📵

Department of Biomechanics and Ergonomics, Faculty of Physical Therapy, Cairo University, Egypt.



Citation Ahmed EA, Nassif NS, Radwan RE, Elborady AA, Mohamed YE. Effects of Hamstring Flossing on Balance and Foot Function in Female Patients with Plantar Fasciopathy: A Randomized Controlled Study. Journal of Modern Rehabilitation. 2025;19(4):405-414. http://dx.doi.org/10.18502/jmr.v19i4.19777

doi http://dx.doi.org/10.18502/jmr.v19i4.19777

Article info:

Received: 23 Mar 2025 Accepted: 03 Jun 2025

Available Online: 01 Oct 2025

ABSTRACT

Introduction: A relationship between hamstring tightness and plantar fasciopathy (PF) has been reported. Hamstring flossing has been reported to improve hamstring flexibility; however, its effects on balance and foot function in patients with PF have not been investigated. This study aimed to compare the efficacy of a conventional physical therapy treatment protocol combined with hamstring flossing versus a conventional physical therapy treatment protocol alone on balance and foot function in patients with PF.

Materials and Methods: Thirty patients with PF were randomly assigned to two groups: The study group (A) received hamstring flossing in addition to the conventional physical therapy treatment protocol, whereas patients in the control group (B) received only the conventional physical therapy treatment protocol. Foot function was evaluated using the foot function index (FFI), and the Biodex balance system (BBS) was used to assess the overall stability index (OSI). Outcomes were evaluated at baseline and three weeks after the intervention. The groups were compared using a two-way mixed-design multivariate analysis of variance (MANOVA).

Results: The post-test mean values of the FFI score and OSI were significantly higher than the pre-test values in both the flossing and control groups (P=0.001). No significant intergroup differences were observed for either variable post-treatment (P>0.05).

Conclusion: The combination of conventional treatment for PF and hamstring flossing provided no additive benefit compared to conventional treatment alone in patients with PF.

Keywords:

Fasciitis; Plantar; Hamstring flossing; Balance; Foot function

* Corresponding Author:

Eman Ahmed Ahmed, PhD.

Address: Department of Biomechanics and Ergonomics, Faculty of Physical Therapy, Cairo University, Egypt.

Tel: +20 (20) (11) 22211657 **E-mail:** eman3ahmed@yahoo.com



Introduction

lantar fasciitis (PF) is a condition that develops gradually due to overuse. It is most frequently observed in both non-athletic and athletic populations [1]. It affects approximately 10% of the world's population at least once in their lifetime, between the ages of 40–60 years [2]. Approximately 80% of all heel pain cases are attributed to PF, with a higher incidence in females [3]. The primary complaint of patients with PF is startup pain, characterized by a sharp discomfort at the medial plantar heel during the first steps of the morning or after rest. Although this pain diminishes with walking, it typically worsens at the end of the day or after high-impact activities [4].

PF is a multifactorial disorder with several mechanical factors contributing to its development. The prevailing theory suggests that chronic biomechanical stress triggers microfascial breakdown, impairs healing, and causes abnormal histological responses [5]. Functional deficits, such as calf weakness, Achilles tendon/plantar fascia tension, gastrocnemius tightness, poor foot alignment, and hamstring tightness, increase strain on the relatively inelastic plantar fascia [6, 7]. The incidence of PF is 8.7-fold higher in patients with hamstring tightness [8].

Standard PF management includes patient education [5] and conservative treatments (e.g. stretching, strengthening, ultrasound, heat therapy, medication, and steroid injections) [9]. A key treatment goal is to restore normal tissue flexibility [10]; hence, stretching of the plantar fascia, calf, and Achilles tendon is recommended. Despite short-term benefits, many patients report persistent pain post-rehabilitation, underscoring the need for better therapies [11].

Given the link between hamstring tightness and PF, techniques, such as static stretching and PNF have been used to improve hamstring flexibility [12, 13]. However, comparative evidence for these methods is limited. Unlike traditional stretching (which compromises strength), flossing may enhance ROM without reducing power. This technique uses a compressive latex band to temporarily restrict blood flow, followed by exercises to break myofascial adhesions [14].

Despite its potential, the effects of hamstring flossing on balance and function in patients with PF remain unknown. Tissue flossing is emerging for improving ROM, pain, and injury prevention. Kaneda et al. [15] found flossing superior to stretching for knee ROM and

eccentric strength. However, no studies have examined hamstring flossing's impact on PF-related balance and function (foot function index [FFI]/overall stability index [OSI]).

We hypothesized that hamstring flossing would significantly improve FFI and OSI in patients with PF. This randomized controlled trial aimed to evaluate the effect of hamstring flossing in combination with conventional therapy versus conventional therapy alone, and to compare changes in FFI and OSI between the two groups.

Materials and Methods

Thirty patients with PF participated in this study. The patients' mean values of age, body mass, height, and body mass index (BMI) were 45.93±8.62 years, 85.26±12.92 kg, 159.66±7.02 cm, and 30.35±4.03 kg/m², respectively. They were randomly assigned to two equal groups using a computer-generated method. Patients in the study group (A) received hamstring flossing exercises in addition to the conventional physical therapy treatment program, which included therapeutic ultrasound, moist heat, Achilles tendon stretching, plantar fascia stretching, and strengthening exercises for the intrinsic foot muscles [16]. The control group (B) received only the conventional physical therapy treatment program. All patients read and signed an informed consent form prior to commencing the study.

Study setting and timeline

This study was conducted at the Physical Therapy Outpatient Clinic of the Faculty of Physical Therapy, Cairo University. The practical aspect of this study spanned 16 months, from October 2021 to January 2023. The inclusion criteria included female patients with PF who had limited hamstring muscle flexibility (20° or more loss of knee extension), as measured by the active knee extension test [17]. Participants also had heel pain and tenderness at the medial tubercle of the calcaneus (the site of insertion of the plantar fascia) with a pain intensity greater than 4 out of 10 on the visual analog scale. Their pain was at its highest level during the first steps in the morning after waking up and during walking after a period of rest. Clinical diagnosis was confirmed using the windlass test; the appearance of pain or increased pain at the insertion of the plantar fascia indicated a positive test for PF [18].

Exclusion criteria

Patients were excluded from the study if they had a history of trauma, strain, sprain, deformity, or radiating pain in the hamstring muscle or plantar area of the foot within the last six months. Additional exclusion criteria included a BMI greater than 35 kg/m², latex allergy, hypertension, systemic inflammatory disease, or venous thrombotic disease [15, 11].

Instrumentation

Biodex balance system (BBS)

The OSI, a quantitative measure of balance, was assessed using the BBS (Biodex Medical Systems, Inc., Shirley, NY). The BBS is a validated and reliable tool for objectively measuring a patient's ability to maintain balance on both stable and unstable surfaces [19]. Higher OSI values indicate poorer balance.

Sphygmomanometer

Floss band compression during treatment was monitored using a modified sphygmomanometer [20] that operates on the same principles as the Kikuhime pressure sensor [21].

3. Voodoo flossing band

A Sanctband COMPRE Floss Blueberry band (5.1 cm × 3.5 m; Sanct Japan Co., Ltd.) was used for hamstring flexibility training. The flossing protocol involved wrapping the latex band around the target tissue for 1–3 minutes [14], maintaining an interface pressure of 140–160 mm Hg (measured using a sphygmomanometer), and guiding the patient to move the flossed joint through its full range of motion [18].

Procedures

The study consisted of four phases: initial assessment, treatment, reassessment, and statistical analysis.

Initial assessment phase

This study used a pre-test-post-test control group design, in which two groups of patients were tested before and after treatment. Balance and foot function were assessed for each patient in both groups.

Balance testing via BBS

The platform of the BBS was set at firmness level eight, the most stable setting. Each patient stood barefoot on the platform with their hands beside their body while maintaining a comfortable position. The BBS screen was adjusted to eye level according to each patient's height, and handrails were positioned for safety in case of a loss of balance. The test began when patients were informed that "the plate is now unlocked." With eyes open, patients were instructed to keep the platform level by centering the cursor on the bull's-eye of the screen. The test duration was preset to 20 seconds. After completion, the platform was automatically locked, and the software averaged the results. The OSI was recorded as the mean of three trials.

FFI

The FFI measured the impact of foot pathology on function in terms of pain, disability, and activity restriction. Patients in both groups underwent testing before and after a three-week intervention. Each patient scored questions on a 0–10 scale (0=no pain/difficulty; 10=worst pain/difficulty). Subscale scores were calculated by dividing the total score by the maximum possible score and multiplying by 100. Scores ranged from 0% (no impairment) to 100% (maximum impairment) [22].

Treatment phase

Hamstring flossing band procedures

A Voodoo floss band (2 m × 5 cm; Mobility WOD) was applied to the hamstring muscle of the affected leg three times a week for three weeks [15]. The patients stood with the affected foot forward. At the same time, the examiner wrapped the band from distal to proximal over the distal third of the thigh, between the anterior superior iliac spine and patella, with each wrap overlapping half of the previous one. The therapist then performed four passive twists of the wrapped band before guiding the patient through two minutes of active knee flexion/extension (to maximum range of motion). The exercise was repeated after a one-minute rest. Finally, the band was removed, and patients walked for one minute to restore circulation [21].

Despite the general safety of flossing, several precautions were implemented. The patients were instructed to report any strong tingling or pain during application, which would require immediate removal of the band. Skin color was checked post-removal by applying and releasing finger pressure to verify normal capillary refill. The application duration was adjusted (2–5 minutes) based on individual tolerance [23]. All participants were advised to maintain their regular activity levels and refrain from participating in other training programs [24, 25].

Conventional physical therapy protocol

Therapeutic ultrasound (continuous mode: 1.5 W/cm², 3 MHz) was applied for a maximum of seven minutes at the most painful plantar site three times weekly for three weeks [16]. For Achilles tendon stretching, patients faced a wall with hands supported, aligned the affected leg behind the non-affected leg, and bent the front knee while keeping the back knee straight to feel a calf/Achilles stretch. This 10-second hold was repeated 15 times twice daily for three weeks (performed both in-clinic and at home). Plantar fascia stretching involved sitting with the affected leg crossed and pulling the toes toward the shin until a stretch was felt in the arch (20-second hold, 15 repetitions twice daily). Intrinsic foot muscle strengthening ("towel toe curls") required patients to grasp a towel with their toes (with a 3-second hold) and performed 100 repetitions daily. Additionally, the patients submerged the affected foot in warm water for 20 minutes nightly.

Re-assessment phase

After three weeks of treatment, all patients were assessed for OSI and FFI scores.

Statistical analysis

Two independent variables were examined: group (flossing + conventional therapy vs. conventional therapy alone) and time (pre- vs post-treatment), with FFI and OSI as dependent variables. The sample size was calculated using G*Power software, version3.1.9.6, assuming a medium effect size (Cohen's d=0.50) and 80% power. Of the 45 initially recruited patients, 30 completed the study, resulting in an overall dropout rate of 33% (21% after enrollment). Normality was confirmed using the Kolmogorov-Smirnov/Shapiro-Wilk tests and distribution analysis. Parametric tests included unpaired t-tests (demographics), chi-square tests (affected-side distribution), Levene's test (homogeneity), and two-way mixeddesign multivariate analysis of variance (MANOVA) (FFI/OSI comparisons. Analyses were performed using SPSS software, version 25 (IBM), with significance at P<0.05.

Results

Flowchart of participants

Of the 45 patients who initially met the inclusion criteria, seven declined to participate. The remaining 38 patients were randomly allocated to two groups. Five patients withdrew after initial evaluation due to work commitments (three from group A and two from group B). Two group B participants dropped out after three weeks due to personal reasons, and one Group A patient was excluded from the analysis. Consequently, 30 female patients with unilateral PF completed the study (Figure 1).

Patients' demographic data

Unpaired t-tests and chi-square tests showed no significant differences between groups for age, weight, height, BMI, or affected side distribution (P>0.05) (Table 1).

Effect of treatment on FFI and OSI

Mixed-design multivariate analysis of variance (MANOVA) revealed significant reductions in FFI scores post-treatment compared to baseline in both groups (P<0.001) (Figure 2). Similarly, OSI values decreased significantly post-treatment (P<0.001) (Figure 3). Between-group comparisons showed no significant differences in pretreatment or post-treatment values for either FFI or OSI (P>0.05) (Table 2).

Discussion

Within-subject effects (conventional treatment with hamstring flossing)

This study investigated the effects of adding hamstring flossing to conventional physical therapy on the FFI and OSI in 30 women with PF. The significant post-treatment improvements in both groups may be attributed to Achilles tendon and plantar fascia stretching, which targets calf muscle tightness, a key contributor to PF [26]. Stretching these structures remains a cornerstone of treatment [27], with Engkananuwat et al. [28] demonstrating that combined Achilles tendon and plantar fascia stretching improves pain and ankle dorsiflexion ROM more effectively than isolated Achilles stretching.

Stretching enhances joint ROM and balance [29], while plantar fascia-specific stretching is critical for arch stability via the windlass mechanism. Restricted ankle dorsiflexion due to Achilles tightness forces compensatory subtalar motion, increasing stress on the plantar




Figure 1. Participants, flow chart

fascia and promoting overpronation, a pathway to fasciopathy and flatfoot [30].

In the current study, the improvement in foot function observed in both groups may be linked to the combined effects of therapeutic ultrasound and moist heat application. Therapeutic ultrasound has been shown to stimulate the healing process [11], while moist heat significantly improves local circulation and reduces pain in the plantar fascia [31]. Sullivan et al. [32] highlighted that weakness in intrinsic and extrinsic foot muscles may contribute to symptom recurrence in PF patients due to impaired muscle function and joint positioning during gait. Consequently, strengthening the intrinsic foot muscles is crucial for counteracting excessive stress on



the plantar fascia [24]. Supporting this, Wei et al. [33] reported that intrinsic foot muscle training enhances biomechanical function of the medial longitudinal arch and improves dynamic postural balance. The intrinsic foot muscle strengthening protocol implemented in this study likely contributed to the observed improvement in heel pain.

The significant post-treatment reduction in OSI and FFI scores in the flossing group may reflect improved hamstring flexibility achieved through flossing band application. Behm et al. [34] demonstrated that floss bands generate compressive stress on target muscles, skin, and fascia, modifying fluid viscosity and reducing movement resistance. Similarly, Kaneda et al. [15] found that




Figure 2. Mean values of the FFI score pre, and 3 weeks post-treatment for patients in both tested groups

flossing induces blood flow restriction and localized vascular occlusion, which alters fascial-neuromuscular interactions to facilitate greater fascial mobility. Notably, Kaneda et al. [15] further established that hamstring flossing surpasses dynamic stretching in improving hamstring flexibility in healthy young males.

Between-subject effect

The current study found no statistically significant differences in the OSI or FFI between the control and experimental groups. This may be explained by the conventional treatment program's inclusion of gastrocnemius muscle stretching, which could have indirectly improved hamstring flexibility [35], the primary target of hamstring flossing.




Figure 3. Mean values of the OSI of balance pre, and 3 weeks post-treatment for patients in both tested groups

JMR

JMR

JMR

Table 1. Comparisons of patients' characteristics between the study and control groups

Patients' Characteristics		Mean±SD/No. (%)			
		Study Group	Control Group	t	Р
Age (y)		47.2±9.7	44.9±9.5	0.67	0.51
Weight (Kg)		82.4±13.8	84.1±15.9	-0.32	0.75
Height (cm)		159.4±7.7	157.1±8.1	0.81	0.42
BMI (Kg/m²)		30.35±4.03	30.35±4.03	-0.89	0.37
Affected side	Right side	7(44)	7(47)	(χ²=0.02)	0.07
	Left side	9(56)	8(53)		0.87

 χ^2 : Chi-squared value; BMI: Body mass index.

Myers [36] conceptualized the superficial back line as a myofascial chain connecting the hamstrings proximally to the gastrocnemius and plantar fascia distally. This anatomical relationship suggests that hamstring flexibility enhances the efficacy of calf muscle stretching, thereby improving foot pain and function. Fauris et al. [37] supported this, demonstrating that flexibility training on any segment of this chain increases hamstring and calf flexibility while improving ankle dorsiflexion range. Although hamstring flexibility training may benefit PF treatment, targeted interventions for the plantar fascia remain equally impactful [38], potentially explaining the comparable outcomes between groups.

Gastrocnemius shortening often manifests as calf pain before hamstring tightness, limiting hamstring flexibility testing due to gastrocnemius-related discomfort [39]. Russell et al. [35] investigated this dynamic, comparing hamstrings-only, gastrocnemius-only, and combined stretching. Their results, consistent with ours, highlighted the unique role of gastrocnemius stretching in improving knee extensibility, likely due to its biarticular anatomy spanning the knee and ankle.

Contrasting opinions

Contrary to our findings, Radford et al. [40] reported that two weeks of calf muscle stretching failed to sig-

Table 2. Mean FFI and OSI pre and post-treatment of study and control groups

Variables		Mean±SD			_	Fff + 6:
		Group A	Group B	MD	Р	Effect Size
FFI	Pre-treatment	0.72±0.07	0.70±0.08	0.02	0.57	
	Post-treatment	0.40±0.15	0.42±0.11	-0.02	0.67	0.15
	MD	0.32	0.28			
	% of change	44.44	40			
		P=0.001	P=0.001			
OSI	Pre-treatment	3.68±0.86	3.71±0.92	-0.03	0.92	
	Post-treatment	2.73±0.76	2.83±1.11	-0.46	0.18	0.48
	MD	1.31	0.88			
	% of change	35.60	23.72			
		P=0.001	P=0.001			

Abbreviations: MD: Mean difference; FFI: OSI: Overall stability index; FFI: Foot function index.



nificantly improve first-step pain, foot pain, or foot function in 92 patients with plantar fasciitis compared to a non-stretching control. This discrepancy may stem from methodological differences: Radford et al. [40] used a wedge-based stretching protocol, which redistributed pressure to the plantar heel, potentially exacerbating discomfort. In contrast, our study employed dynamic lunge stretches for the Achilles tendon, a method associated with improved muscle strength [41]. The strength gains from dynamic stretching may have enhanced extrinsic foot muscle function, compensating for the impaired arch stabilization and reduced ground reaction force absorption characteristics of PF [42].

Our results align with Wu et al. [43], who demonstrated that hamstring flossing enhanced knee ROM without compromising static balance in healthy women. This supports the premise that flossing improves flexibility without negatively affecting postural control, a critical consideration for PF rehabilitation.

Conclusion

The addition of hamstring flossing to conventional physical therapy did not significantly enhance foot function or balance in patients with PF compared to conventional treatment alone.

Study limitations

Lack of a non-intervention control group, which limits our ability to assess spontaneous symptom progression.

Variable flossing band pressure due to subjective discomfort and differences in application technique.

Limited generalizability beyond the studied age group and sex (female patients aged 40–60 years).

Exclusion of balance training, whose potential effects on functional improvement remain unexplored.

Short intervention duration (3 weeks), which may not capture long-term outcomes.

Future directions

Placebo-controlled trials to isolate the effects of hamstring flossing.

Inclusion of a non-intervention group to benchmark natural symptom progression.

Integration of neuromuscular training for ankle/foot muscles and long-term follow-up.

Investigation of balance training as an adjunct therapy for PF.

Ethical Considerations

Compliance with ethical guidelines

This study adhered to the ethical standards outlined in the Declaration of Helsinki and was approved by the Research Ethics Committee of the Faculty of Physical Therapy, Cairo University, Cairo, Egypt (Code: P.T.REC/012/004739). The trial was approved by the Pan African Clinical Trial Registry, Cape Town, South Africa (Code: PACTR202408921930700). The study was retrospectively registered on August 8, 2024.

Funding

This research did not receive any grant from funding agencies in the public, commercial, or non-profit sectors.

Authors' contributions

Conceptualization and supervision: Amal Abdelrahman Elborady and Yassmin Essam Mohamed; Methodology: Eman Ahmed Ahmed, Nagui Sobhi Nassif, and Rafik Elmaamoon Radwan; Data collection and analysis: Eman Ahmed Ahmed; Investigation and writing: Eman Ahmed Ahmed, Amal Abdelrahman Elborady, and Yassmin Essam Mohamed.

Conflict of interest

The authors declared no conflict of interest.

Acknowledgments

The authors thank all the participants in the study for their cooperation.

References

- [1] Salehi S, Shadmehr A, Olyaee G, Bashardoust Tajali S, Mir S M. Effectiveness of dry needling for the management of plantar fasciitis: A review study. Journal of Modern Rehabilitatio. 2019; 13(1):1-10. [DOI:10.32598/jmr.13.1.1]
- [2] Rhim HC, Kwon J, Park J, Borg-Stein J, Tenforde AS. A systematic review of systematic reviews on the epidemiology, evaluation, and treatment of plantar fasciitis. Life. 2021; 11(12):1287. [DOI:10.3390/life11121287] [PMID]

- [3] Palomo-López P, Becerro-de-Bengoa-Vallejo R, Losa-Iglesias ME, Rodríguez-Sanz D, Calvo-Lobo C, López-López D. Impact of plantar fasciitis on the quality of life of male and female patients according to the foot health status questionnaire. Journal of Pain Research. 2018; 11:875-80. [DOI:10.2147/JPR.S159918] [PMID]
- [4] Iglesias M, Sperone E, Macklin Vadell A, Bigatti A. Plantar fasciitis: Analysis of therapeutic options. Revista de la Asociación Argentina de Ortopedia y Traumatologia. 2022; 87(3):413-21. [DOI:10.15417/issn.1852-7434.2022.87.3.1359]
- [5] Mushtaq N, Abbasian A. Plantar fasciitis. Annals of the Royal College of Surgeons of England. 2013; 95(3):229. [DOI:10.13 08/003588413X13511609957812] [PMID]
- [6] Shashua A, Flechter S, Avidan L, Melayev A, Kalichman L. The effect of additional ankle and midfoot mobilizations on plantar fasciitis: A randomized controlled trial. The Journal of Orthopaedic and Sports Physical Therapy. 2015; 45(4):265-72. [DOI:10.2519/jospt.2015.5155] [PMID]
- [7] Tariq H, Mazhar F, Ghaffar E, Arif T, Irshad U, Aqdas A. Association between plantar fasciitis and hamstring tightness among runners. Journal of Health and Rehabilitation Research. 2024; 4(2):343-7. [DOI:10.61919/jhrr.v4i2.803]
- [8] Sayed MA, El-megid SF, El-azizy HM, El-Khozamy HM. The correlation between hamstring tightness and plantar fasciitis. The Medical Journal of Cairo University. 2019; 87(1):309-13. [DOI:10.21608/mjcu.2019.52353]
- [9] Manju, Meetu, Pradyumna Krishna, Sameer. Effect of various conventional treatments on plantar fasciitis: A brief review. International Journal of Science and Healthcare Research. 2023; 8(3):276-80. [DOI:10.52403/ijshr.20230338]
- [10] Ismail MM, Ahmed S. The Incorporation of balancing board exercises in the management of plantar fasciitis. International Journal of Health Sciences. 2011; 16(1):19-28. [Link]
- [11] Latt LD, Jaffe DE, Tang Y, Taljanovic MS. Evaluation and treatment of chronic plantar fasciitis. Foot & Ankle Orthopaedics. 2020; 5(1):2473011419896763. [DOI:10.1177/2473011419896763] [PMID]
- [12] Vidhi S, Anuprita T, Asmita K, et al. Comparison of PNF technique with NDS technique for hamstrings tightness in asymptomatic subjects. The Indian Journal of Physiotherapy & Occupational Therapy. 2014; 8(3):158-61. [DOI:10.5958/0973-5674.2014.00375.X]
- [13] Ganacharya PM, Pagare VK, Sareen A, Palekar TJ. Effect of neurodynamic sliding technique versus static stretching on hamstring flexibility in football players with short hamstring syndrome. Journal of Musculoskeletal Research. 2014; 17(2):1751-63. [DOI:10.1142/S0218957714500092]
- [14] Maust Z, Bradney D, Collins SM, Wesley C, Bowman TG. The effects of soft tissue flossing on hamstring range of motion and lower extremity power. International Journal of Sports Physical Therapy. 2021; 16(3):689-94. [DOI:10.26603/001c.24144] [PMID]
- [15] Kaneda H, Takahira N, Tsuda K, Tozaki K, Kudo S, Takahashi Y, Sasaki S, Kenmoku T. Effects of tissue flossing and dynamic stretching on hamstring muscles function. Journal of Sports Science & Medicine. 2020; 19(4):681. [PMID]

- [16] Prakash S, Anand M. Effect of manual therapy versus conventional therapy in patients w ith plantar fasciitis—A comparative study. International Journal of Physiotherapy and Research. 2014; 2(1):378-82. [Link]
- [17] Bolívar YA, Munuera PV, Padillo JP. Relationship between tightness of the posterior muscles of the lower limb and plantar fasciitis. Foot & Ankle International. 2013; 34(1):42-8. [DOI:10.1177/1071100712459173] [PMID]
- [18] Garceau De D, Dean D, Requejo SM. The association between diagnosis of plantar fasciitis and windlass test results. Foot & Ankle International. 2003; 24(3):251-55. [DOI:10.1177/107110070302400309] [PMID]
- [19] Karimi N, Ebrahimi I, Kahrizi S, Torkaman G. Evaluation of postural balance using the biodex balance system in subjects with and without low back pain. Pakistan Journal of Medical Sciences. 2008; 24(3):372. [Link]
- [20] Konrad A, Bernsteiner D, Budini F, Reiner MM, Glashüttner C, Berger C, et al. Tissue flossing of the thigh increases isometric strength acutely but has no effects on flexibility or jump height. European Journal of Sport Science. 2021; 21(12):1648-58. [DOI:10.1080/17461391.2020.1853818] [PMID]
- [21] Driller M, Mackay K, Mills B, Tavares F. Tissue flossing on ankle range of motion, jump and sprint performance: A follow-up study. Physical Therapy in Sport. 2017; 28:29-33. [DOI:10.1016/j.ptsp.2017.08.081] [PMID]
- [22] Rutkowski R, Gizińska M, Gałczyńska-Rusin M, Kasprzak MP, Budiman-Mak E. The importance of foot function assessment using the foot function index-revised short form (FFI-RS) questionnaire in the comprehensive treatment of patients with rheumatoid arthritis. Journal of Clinical Medicine. 2022; 11(9):2298. [DOI:10.3390/jcm11092298] [PMID]
- [23] Kim J, Sung DJ, Lee J. The therapeutic effect of the flossing technique on knee flexibility and functional performance: A pilot study. Journal of Sports Science and Medicine. 2017; 16:130-6. [Link]
- [24] Huffer D, Hing W, Newton R, Clair M. Strength training for plantar fasciitis and the intrinsic foot musculature: A systematic review. Physical Therapy in Sport. 2017; 24:44-52. [DOI:10.1016/j.ptsp.2016.08.008] [PMID]
- [25] Cohen J. Statistical power analysis for the behavioral sciences. 2nd ed. New York: Routledge; 1988. [DOI:10.4324/9780203771587]
- [26] Pearce CJ, Seow D, Lau BP. Correlation Between Gastrocnemius Tightness and Heel Pain Severity in Plantar Fasciitis. Foot & Ankle International. 2021; 42(1):76-82. [DOI:10.1177/1071100720955144] [PMID]
- [27] Gupta R, Malhotra A, Masih GD, Khanna T, Kaur H, Gupta P, et al. Comparing the role of different treatment modalities for plantar fasciitis: A double blind randomized controlled trial. Indian Journal of Orthopaedics. 2020; 54(1):31-37. [DOI:10.1007/s43465-019-00038-w] [PMID]
- [28] Engkananuwat P, Kanlayanaphotporn R, Purepong N. Effectiveness of the simultaneous stretching of the achilles tendon and plantar fascia in individuals with plantar fasciitis. Foot & Ankle International. 2018; 39(1):75-82. [DOI:10.1177/1071100717732762] [PMID]

- [29] Gillespie LD, Robertson MC, Gillespie WJ, Sherrington C, Gates S, Clemson LM, et al. Interventions for preventing falls in older people living in the community. The Cochrane Database of Systematic Reviews. 2012; 2012(9):CD007146. [DOI:10.1002/14651858.CD007146.pub3] [PMID]
- [30] Scattone Silva R, Smitheman HP, Smith AK, Silbernagel KG. Are static foot posture and ankle dorsiflexion range of motion associated with Achilles tendinopathy? A crosssectional study. Brazilian Journal of Physical Therapy. 2022; 26(6):100466. [DOI:10.1016/j.bjpt.2022.100466] [PMID]
- [31] Petrofsky J, Laymon M, Lee H. Local heating of trigger points reduces neck and plantar fascia pain. Journal of Back and Musculoskeletal Rehabilitation. 2020; 33(1):21-8. [DOI:10.3233/BMR-181222] [PMID]
- [32] Sullivan J, Burns J, Adams R, Pappas E, Crosbie J. Musculoskeletal and activity-related factors associated with plantar heel pain. Foot & Ankle International. 2015; 36(1):37-45. [DOI:10.1177/1071100714551021] [PMID]
- [33] Wei Z, Zeng Z, Liu M, Wang L. Effect of intrinsic foot muscles training on foot function and dynamic postural balance: A systematic review and meta-analysis. Plos One. 2022; 17(4):e0266525. [DOI:10.1371/journal.pone.0266525] [PMID]
- [34] Behm DG, Wilke J. Do self-myofascial release devices release myofascia? Rolling mechanisms: A narrative review. Sports Medicine. 2019; 49(8):1173-81. [DOI:10.1007/s40279-019-01149-y] [PMID]
- [35] Russell PJ, Decoster LC, Enea D. Effects of gastrocnemius, hamstring, and combined stretching programs on knee extensibility. Athletic Training & Sports Health Care. 2010; 2(2):67-73. [DOI:10.3928/19425864-20100226-06]
- [36] Myers TW. Anatomy trains: Myofascial meridians for manual and movement therapists. Amsterdam: Elsevier Health Sciences; 2009. [Link]
- [37] Fauris P, López-de-Celis C, Canet-Vintró M, Martin JC, Llurda-Almuzara L, Rodríguez-Sanz J, et al. Does self-myofascial release cause a remote hamstring stretching effect based on myofascial chains? A randomized controlled trial. International Journal of Environmental Research and Public Health. 2021; 18(23):12356. [DOI:10.3390/ijerph182312356] [PMID]
- [38] Siriphorn A, Eksakulkla S. Calf stretching and plantar fascia-specific stretching for plantar fasciitis: A systematic review and meta-analysis. Journal of Bodywork and Movement Therapies. 2020; 24(4):222-32. [DOI:10.1016/j.jbmt.2020.06.013] [PMID]
- [39] Kawano MM, Ambar G, Oliveira BI, Boer MC, Cardoso AP, Cardoso JR. Influence of the gastrocnemius muscle on the sit-and-reach test assessed by angular kinematic analysis. Revista Brasileira de Fisioterapia. 2010; 14(1):10-5. [DOI:10.1590/ S1413-35552010000100003] [PMID]
- [40] Radford JA, Landorf KB, Buchbinder R, Cook C. Effectiveness of calf muscle stretching for the short-term treatment of plantar heel pain: A randomised trial. BMC Musculoskeletal Disorders. 2007; 8:36. [DOI:10.1186/1471-2474-8-36] [PMID]
- [41] Boonchum H, Bovonsunthonchai S, Komsak S, Kunanusornchai W. Effect of a home-based stretching exercise on multisegmental foot motion and clinical outcomes in patients with plantar fasciitis. Journal of Musculoskeletal & Neuronal Interactions. 2020; 20(3):411-20. [PMID]

- [42] Lee J, Kim J. Effects of an 8-week lunge exercise on an unstable support surface on lower-extremity muscle function and balance in middle-aged women. Physical Activity and Nutrition. 2022; 26(4):14-21. [DOI:10.20463/pan.2022.0020]
- [43] Wu SY, Tsai YH, Wang YT, Chang WD, Lee CL, Kuo CA, et al. Acute effects of tissue flossing coupled with functional movements on knee range of motion, static balance, in single-leg hop distance, and landing stabilization performance in female college students. International Journal of Environmental Research and Public Health. 2022; 19(3):1427. [DOI:10.3390/ijerph19031427] [PMID]