Review Article

Comparison of Functional Movement Screen Scores and Performance Between Male and Female Soccer Players

Alireza Zare[1], Hashem Piri[2*], Razieh Hajizadeh[1], Amir Reza Saeidizadeh[1], Amir Reza Ghareh Khanlou[1]

[1]- Department of Corrective Exercise and Sport Injury, Faculty of Physical Education and Sport Sciences, Science and Research Branch, Islamic Azad University, Tehran, Iran
[2]- Department of Corrective Exercise and Sport Injury, Faculty of Physical Education and Sport Sciences, Allameh Tabataba'i University, Tehran, Iran

*** Corresponding author:** *Hashem Piri, Assistant Professor*
Department of Corrective Exercise and Sport Injury, Faculty of Physical Education and Sport Sciences, Allameh Tabataba'i University, Tehran, Iran
Emails: hpiri1984@gmail.com

ORCID ID:
Alireza Zare: 0009-0009-6919-7271
Hashem Piri: 0000-0002-3137-6161
Razieh Hajizadeh: 0009-0001-8719-1879
Amir Reza Saeidizadeh: 0009-0009-0664-218X
Amir Reza Ghareh Khanlou: 0009-0002-9835-3631

Article info:
Received: 13 May 2025
Accepted: 7 Jun 2025

Citation: Zare A, Piri H, Hajizadeh R, Saeidizadeh AR, Ghareh Khanlou AR. Comparison of Functional Movement Screen Scores and Performance Between Male and Female Soccer Players. Journal of Modern Rehabilitation. 2026;20(1):?-?

Running Title: FMS Scores & Performance in Male vs Female Soccer Players

Abstract:
Introduction: Soccer is the most popular sport globally for both men and women. The Functional Movement Screen (FMS) is a seven-part standardized tool created by experts to evaluate movement quality. This research compared FMS scores and performance between male and female soccer players.
Materials and Methods: A total of sixty semi-professional soccer players (30 males and 30 females) participated in this study. All participants completed the FMS test, Y-Balance Test, Davies Test, Single hop for distance, and Crossover Hop Test. Between-group comparisons were performed using independent t-tests for normally distributed variables and Mann-Whitney U tests for non-normally distributed variables.

Results: There were no significant differences between male and female players in Deep Squat, Hurdle Step, In-Line Lunge, or Overall FMS Score (P>0.05). However, significant differences were found in Shoulder Mobility, Active Straight Leg Raise, Trunk Stability Push-Up, and rotary stability (P<0.05). Males performed better in Trunk Stability, while females excelled in Shoulder Mobility and Active Straight Leg Raise. Males outperformed females in Single hop for distance and Cross-over hop. In upper extremity performance, males performed better in the Davies test and dynamic balance in the superolateral direction, with no significant differences in other upper extremity dynamic balance directions.
Conclusion: This study demonstrates the necessity of gender-specific training programs, as they are crucial in enhancing performance and reducing the risk of injury. According to the findings, it is recommended that flexibility and mobility exercises be implemented for male soccer players and core stability and strength training for female soccer players.
Keywords: Soccer, Athletic Performance, Risk Factors, Sex.

1. Introduction

Soccer is widely regarded as one of the most prominent sports worldwide, attracting the largest global audience. It promotes both physical and psychosocial development and demands high levels of strength, endurance, agility, speed, flexibility, and balance (1, 2). In addition, soccer is classified as a sport with a high incidence of injury, primarily due to the unpredictable nature of complex field situations, the intensity of play, and the inherent physical contact between players (3). As a result, at the professional level, injury prevention and minimizing player absence during training and matches are critical. Given the high financial investment in player contracts and tournament participation, preventing and managing sports injuries are very important (4). Previous studies have emphasized the role of proper movement patterns in ensuring safe and efficient long-term physical performance in soccer players (5-7).

Optimal movement patterns and strong neuromuscular control are essential for improving performance and reducing injury risk, particularly during high-intensity actions such as landing, cutting, and rapid directional changes common in soccer (٨). Since 2004, the number of athletes engaged in competitive sports has increased significantly, with greater focus on athlete safety. These has led to extensive research in injury prevention through effective screening and rehabilitation strategies. One widely recognized tool in this context is the FMS, which aims to detect and correct dysfunctional or high-risk movement patterns that may predispose athletes to injury (9). Introduced by Cook, Burton, and Hoogenboom in 2006 (10), the FMS is a practical assessment tool designed to identify asymmetries and limitations in mobility and stability. Its primary goal is to detect movement dysfunctions that may elevate the risk of injury.

The FMS evaluates seven fundamental movement patterns that assess balance, mobility, and stability, reflecting key principles of proprioception and kinesthetic awareness .(١٠) A composite score from all seven tasks helps identify limitations in flexibility, bilateral strength asymmetries, and core stability deficiencies factors closely related to athletic performance potential and injury risk .(١١) Moreover, core muscle weakness is widely recognized as a contributor to injury, as core stability is essential for efficient force transfer between the upper and lower body. Weakness in the hip and thigh muscles can lead to instability during lower limb movements, which may result in poor movement patterns and increased injury risk. The FMS helps prevent such issues by detecting them early (12).

Several recent studies have investigated the relationship between FMS scores, physical performance, and gender in soccer players. In the study by Martín-Moya et al. (2023), the

researchers examined the differences and relationships between Functional Movement Screen scores and physical fitness tests in semi-professional male and female soccer players. The findings revealed that male players performed significantly better than female players in the 505 change of direction test. However, no significant gender differences were found in total FMS scores .(۱۳) In contrast, Amjad et al. (2023) found that female footballers scored lower on the composite FMS score compared to male footballers (14).

Evaluating physical performance provides valuable insights into an athlete's physical capabilities (۱۵) and is a fundamental component of elite soccer player development programs (16). In this regard, Mascherini et al. (2024) demonstrated that male athletes excelled in all physical performance tests, including the horizontal jump, triple jump, and 5m and 10m sprints. This superior performance, especially in tests requiring higher power and speed, is likely attributed to greater muscle mass and lower body fat percentage in males (17). Moreover, Barin et al. (2020) conducted a comparative study on the performance of male and female soccer players. The findings revealed that male athletes outperformed their female counterparts in several individual technical skills including vertical jump height, sprint speed, agility, and dribbling power. These performance disparities were partially attributed to inherent differences in anthropometric characteristics between the sexes (18). The data can guide coaches in talent identification, player selection, and long-term athlete development strategies (19).

Although tools such as the FMS are widely utilized, few studies have thoroughly examined individual FMS components and functional performance assessments involving both upper and lower limbs. These assessments are essential for identifying muscular imbalances, postural misalignments, and joint mobility restrictions that can influence performance and injury risk. This gap is particularly relevant given the physiological and biomechanical differences between sexes. As such, comparing FMS scores and functional performance outcomes across genders may offer valuable insights for optimizing training strategies and minimizing injury risk. Accordingly, the present study aimed to compare FMS scores and physical performance between male and female soccer players and to determine whether sex-related differences exist in specific functional performance tasks.

2. Materials and Methods

2.1. Participants

This research was descriptive-comparative and employed a field-based data collection approach.

The study population was young male and female athletes participating in the Tehran Football League. A total of sixty semi-professional soccer players were recruited for this study: thirty males (age: 19.83 ± 1.48 years; weight: 72.40 ± 3.69 kg; height: 176.66 ± 5.24 cm; BMI: 20.30 ± 1.70 kg/m²) and thirty females (age: 19.73 ± 1.48 years; weight: 58.96 ± 5.64 kg; height: 164.36 ± 4.93 cm; BMI: 19.43 ± 1.79 kg/m²). Participants were selected according to predefined inclusion and exclusion criteria. The inclusion criteria were as follows: Participants aged 18-24 years with at least 3 years of membership in a Tehran League team, no significant skeletal or muscular injuries, and no history of surgery in the past 6 months. Participants were excluded if they sustained an injury during the study, withdrew from participation, or failed to complete the research process.

The sample size, calculated using G*Power (version 3.1) (with an effect size of 0.85, $\alpha = 0.05$, and power = 0.8), was 60 participants. The study was approved by the Sport Sciences Research Institute of Iran (Approval Code: SSRI.REC-2310-2493).

2.2. Procedures

Participants were recruited through visits to men's and women's soccer clubs across Tehran. All participants provided written informed consent and completed a demographic information questionnaire before collecting data. Functional movement quality was assessed using the FMS (Functional Movement Screen) with a standardized FMS kit. The Upper Quarter Y-Balance Test (UQ-YBT) evaluated dynamic upper limb balance. Upper limb performance was assessed through the Davies Test, while lower limb performance was measured using the Single hop for distance test and the Cross-over hop for distance Test.

2.3. Measurement

2.3.1. FMS Assessment Protocol

The FMS is an assessment tool to evaluate an individual's fundamental movement patterns. It provides a distinct approach for identifying dysfunctions, predicting performance, and reducing injury risk. Limitations in joint mobility and stability can increase the likelihood of injury among athletes and the general population, whereas optimal movement patterns help prevent and minimize such risks. The FMS consists of seven fundamental movement tasks that require a balance between mobility and stability. These tasks are structured to offer observable performance indicators across basic movement functions such as locomotion, manipulation, and stabilization. By placing athletes in challenging positions where appropriate stability is essential for sufficient mobility, the FMS helps expose weaknesses, asymmetries, and compensatory movement patterns (10, 20).

The FMS comprises seven distinct movement tests: deep squat, hurdle step, in-line lunge, shoulder mobility, active straight-leg raise, trunk stability push-up, and rotary stability. Before the assessment, participants received detailed verbal instructions and demonstrations for each movement pattern. They were allowed one practice trial to familiarize themselves with the movements. During the assessment, participants performed each movement three times. For unilateral tests, the highest score from the three trials was recorded for each side, and the lower of the two scores was used as the final score for that movement. For bilateral tests, the best score from the three trials was recorded (20) (Figure 1).

Evaluators positioned themselves at consistent distances from the participants in anterior, posterior, and lateral views to ensure accurate observation and scoring. Movements were scored on a standardized 4-point scale ranging from 0 to 3. A score of 3 indicates that the movement was performed correctly without any compensations. A score of 2 is given if the movement was completed but with some form of compensation or deviation from the standard movement pattern. A score of 1 is assigned when the individual cannot complete the movement or cannot assume the required position. Finally, A score of 0 is given if pain is reported during any part of the movement (٢٠).

Furthermore, FMS is a reliable assessment tool for interrater (ICC: 0.81; 95% CI: 0.74–0.8) and interrater reliability (ICC: 0.81, 95% CI: 0.70–0.92) (21).

Figure 1. The Functinal Movement Screen Test.

2.3.2. Measurement of Functional Performance

Jump tests were utilized to assess functional performance. A measuring tape was used to determine the distance covered by participants in the zigzag triple jump and single-leg forward hop tests. Additionally, the length of the upper limb was measured from the C7 vertebra to the tip of the middle finger.

2.3.2.1. The Davis Push-Up Test

The Davis Push-Up Test was used to assess upper extremity performance, specifically evaluating muscular strength, endurance, and the stability of the upper limb's closed kinetic chain. This modified push-up test was developed by Davies and Dickoff-Hoffman in 1993 and has demonstrated a high test-retest reliability coefficient of 0.92 (22).

In this assessment, two tape strips were placed on the floor, positioned 90 centimeters (approximately 36 inches) apart. Participants assumed a push-up position between the strips. Over a 15-second interval, they rapidly alternated hand touches to the outer edges of the tape strips in a crisscross manner. The total number of touches completed within this period was recorded as the participant's score. The test was conducted thrice, with the final score calculated as the average of the three trials (22). Moreover, a rest period of 30 to 40 seconds was provided between trials (Figure 2).

Figure 2. Davis Push-Up Test.

2.3.2.2. Single hop for distance test

Single-legged hop tests assess the performance of the entire lower extremity and athletes' ability to perform components of sport-specific actions such as hopping .(٢٣) The single hop for distance test requires a 3-meter-long narrow measuring tape placed on the floor. The athlete stands on their dominant leg, with the toe tip precisely at the tape's starting point. The test involves performing a forward hop to cover the maximum possible distance, landing on the same leg, and maintaining balance for at least 2 seconds. Participants may use their arms for balance if necessary. The participant performed three consecutive single-leg hops using the dominant leg, with the maximum distance achieved recorded as the test result (24, 25). A rest period of 30 to 40 seconds was provided between trials to minimize fatigue. The single hop for distance demonstrated excellent reliability, with an ICC of 0.92 (26) (Figure 3).

2.3.2.3. Cross-over hop for distance test

In this test, the participant stands on the test leg and performs a hopping sequence in a Cross pattern, covering a 15-centimeter-wide line three times. The maximum distance achieved across three trials is recorded as the test outcome. A rest interval of 30 to 40 seconds is provided between each trial to mitigate fatigue and ensure optimal performance.

The total distance covered during the hopping sequence is then recorded. If the participant places the opposite foot on the ground or fails to hop completely over the width of the tape, the trial is considered invalid and must be repeated (27, 28). The cross-over hop for distance showed good reliability, with an ICC of 0.84 (26) **(** Figure 4).



Figure 3. Single hop for distance test **Figure 4.** Cross-over hop for distance

2.3.2.4. Upper Quarter Y-Balance Test (UQ-YBT)

The Upper Quarter Y-Balance Test is a clinical assessment tool used to evaluate upper limb performance in three directions: medial, superolateral, and inferolateral, each at a 135° angle to the other. This test assesses strength, stability, and mobility of the upper extremity and is considered a reliable and valid instrument, with reliability coefficients ranging from 80% to 99% (29). During the test administration, participants shifted their body weight onto the testing arm while maintaining balance on one hand without compromising stability. Simultaneously, they reached as far as possible with the non-testing arm in the three designated directions. The feet were positioned 30 cm apart, and each direction was tested thrice. The average of the three trials in each direction was calculated for analysis, with a two-minute rest interval provided between attempts to reduce fatigue. Given the significant correlation between reach distance and arm length, dynamic balance scores were normalized by dividing the average reach distance in each direction by the participant's arm length and multiplying the result by 100, thereby expressing the reach distance as a percentage of arm length. The composite score was calculated as the sum of the three normalized directional scores (29, 30) **(**Figure 5).

Figure 5. Upper Quarter Y-Balance Test

2.4. Ethical Considerations

All participants were assured that their personal information would remain confidential and accessible only to the research team. They were informed of their right to withdraw from the study at any stage without any repercussions. All procedures were carried out by the ethical standards established by the Sport Sciences Research Institute of Iran. Before the initiation of the study and data collection, participants were provided with detailed information regarding the study's objectives and procedures. Written informed consent was obtained from all participants, indicating their voluntary agreement to take part. Transparency in the research process, the confidentiality of personal data, and the absence of financial obligations for participants were emphasized as fundamental ethical principles maintained throughout the study.

2.5. Statistical analysis

Following data collection, the study participants' dataset was analyzed using descriptive and inferential statistics with SPSS software (version 22). The normality of the data distribution was assessed using the Shapiro-Wilk test. The independent samples t-test was employed for variables with a normal distribution for between-group comparisons. At the same time, the Mann-Whitney

U test was applied for variables with a non-normal distribution. A significance level of 0.05 ($\alpha \leq 0.05$) was set for all statistical tests.

3. Results

Table 1 displays the demographic details of the participants in the study, including age, height, weight, body Mass Index, sports Experience, weekly physical activity, and dominant upper limb along with the mean and standard deviation of the variables and the results of independent T tests. Since the data did not meet the assumption of normality, the Mann–Whitney U test was used to compare the two groups. The results indicated that all variables except for age and sports experience showed statistically significant differences between the groups.

Table 1. The demographic characteristics of the study participants

Variables	**p-value**			**(Mean ± SD) men**	**(Mean ± SD) women**
Age (years)	0.791	(Mean± deviation)	Standard	19.83 ± 1.48	19.73 ± 1.48
Weight (kg)	*0.001	(Mean± deviation)	Standard	72.40 ± 3.69	58.96 ± 5.64
Height (cm)	*0.001	(Mean± deviation)	Standard	176.66 ± 5.24	164.36 ± 4.93
Body Mass Index (BMI)	*0.046	(Mean± deviation)	Standard	20.30 ± 1.70	19.43 ± 1.79
Sports Experience (years)	0.194	(Mean± deviation)	Standard	5.33 ± 1.49	4.83 ± 1.14
Weekly Physical Activity (hours)	*0.001	(Mean± deviation)	Standard	6.60 ± 1.37	5.23 ± 1.13
Dominant Upper Limb Length (cm)	*0.001	(Mean± deviation)	Standard	90.36 ± 4.07	80.56 ± 3.38

Notes: Statistical significance was considered at $p < 0.05$.

The Shapiro-Wilk test was used to assess the normality of the data distribution for both FMS scores and performance (Table 2).

Table 2. Assessment of data normality in research variables

Variable	**(Mean ± SD) men**	**p-value (men)**	**(Mean ± SD) women**	**p-value(women)**

Deep Squat	2.23 ± 0.67	0.001	1.96 ± 0.61	0.001
Hurdle step	2.13 ± 0.57	0.001	1.86 ± 0.57	0.001
In-line lunge	2.23 ± 0.72	0.001	2.16 ± 0.69	0.001
Shoulder Mobility	2.00 ± 0.45	0.001	2.53 ± 0.50	0.001
Active Straight Leg Raise	1.93 ± 0.58	0.001	2.30 ± 0.46	0.001
Trunk Stability Push-up	2.70 ± 0.46	0.001	1.83 ± 0.59	0.001
Rotary stability	2.16 ± 0.46	0.001	1.90 ± 0.54	0.001
Total FMS	15.40 ± 1.90	0.073	14.56 ± 1.83	0.052
Single hop for distance test (m)	1.72 ± 0.18	0.030	1.25 ± 0.15	0.826
Cross-over hop for distance (m)	4.46 ± 0.65	0.224	3.05 ± 0.63	0.115
Davis Test	28.80 ± 5.44	0.132	24.03 ± 3.75	0.199
Dynamic Balance – Medial-Lateral (cm)	97.50 ± 13.19	0.078	94.96 ± 8.82	0.474
Dynamic Balance – Inferior-Lateral (cm)	99.24 ± 17.91	0.720	91.64 ± 12.33	0.248
Dynamic Balance – Superior-Lateral (cm)	92.49 ± 13.97	0.768	84.87 ± 11.76	0.475

Notes: Statistical significance was considered at $p < 0.05$.

According to the results, the FMS subtests and the Single hop for distance test t data was not normally distributed ($P < 0.05$). However, the distribution of data for the total FMS score, the Cross-over hop for distance test, the Davies test, and the UQ-YBT tests in different directions was normal ($P > 0.05$). Accordingly, the Mann-Whitney U test was used to compare groups for non-normally distributed variables. At the same time, the independent t-test was applied to normally distributed variables, at a significance level of 95%.

The results of the between-group comparisons indicated that there were no statistically significant differences between male and female football players in the Deep Squat ($P = 0.108$), Hurdle Step ($P = 0.076$), In-Line Lunge ($P = 0.688$), and Overall FMS Score ($P = 0.089$), as all P-values exceeded the 0.05 threshold. Conversely, significant sex-based differences were observed in Shoulder Mobility ($P = 0.001$), Active Straight Leg Raise ($P = 0.013$), Trunk Stability Push-Up ($P = 0.001$), and Rotary stability ($P = 0.048$). In particular, male football players performed better in the Trunk Stability Push-Up and Rotary stability test. In contrast, female football players outperformed their male counterparts in the Shoulder Mobility and Straight Leg Raise tests. Comparing lower extremity performance between male and female

soccer players revealed significant differences in the Single hop for distance (P = 0.001) and the Cross-over hop for distance (P = 0.001).

Analysis of the mean scores indicated that male soccer players performed better than female soccer players in both the Single hop for distance test and the Cross-over hop for distance tests. Furthermore, comparing upper extremity performance between male and female soccer players showed significant differences in the Davies test (P = 0.001) and UQ-YBT in the superolateral direction (P = 0.026). However, no significant differences were observed between the two groups in the medial (P = 0.384) and inferolateral (P = 0.061) directions of upper extremity dynamic balance. Examination of the mean scores further revealed that male soccer players outperformed female soccer players in the Davies test and upper extremity dynamic balance in the superolateral direction (Table 3 and 4).

Table 3. The Mann-Whitney U test results

Variable	(Mean Rank) Men	(Mean Rank) Women	Z	P-value
Deep Squat	33.73	27.27	-1.609	0.108
Hurdle step	33.83	27.17	-1.774	0.076
Inline lunge	31.33	29.67	-0.402	0.688
Shoulder Mobility	23.30	37.70	-3.779	*0.001
Active Straight Leg Raise	25.90	35.10	-2.493	*0.013
Trunk Stability Push-Up	40.70	20.30	-4.959	*0.001
Rotary stability	33.95	27.05	-1.974	*0.048
Single hop for distance test (m)	1.72	1.25	10.814	*0.001

Table 4. The **T-test Results**

Variable	T	Mean difference	Effect size	P-value
Total FMS score	1.727	0.84	0.049	0.089
Cross-over hop for distance (m)	8.526	1.44	0.556	*0.001
Davis Test	3.946	4.77	0.212	*0.001

Dynamic Balance-Medial-Lateral (cm)	0.876	2.54	0.013	0.384
DynamicBalance - Inferior-Lateral (cm)	1.914	7.60	0.059	0.061
DynamicBalance - Superior-Lateral (cm)	2.287	7.62	0.083	*0.026

4. Discussion

The present study aimed to compare FMS and performance between male and female soccer players. The FMS is recognized as a cost-effective, non-invasive, and reliable tool for evaluating fundamental movement patterns (31, 32). Existing literature supports its high reliability (33, 34) and specificity in identifying athletes at an increased risk of sports-related injuries .(٥) In this study, The FMS was utilized to evaluate fundamental movement patterns. Additionally, upper extremity dynamic balance was measured using the UQ-YBT, upper extremity performance was assessed via the Davies Test, and lower extremity performance was evaluated through the Single hop for distance and Cross-over hop for distance test.

The findings of this study showed that there were no statistically significant differences between male and female football players in several FMS components, specifically in movements such as the Deep Squat, Hurdle Step, In-Line Lunge, and the total FMS score. This suggests that fundamental movement patterns may be similar between genders, particularly when skill level and training are comparable across groups. However, significant gender-related differences were observed in certain FMS components. Specifically, female players performed better in flexibility tests, such as the Shoulder Mobility and Active Straight Leg Raise, which may reflect greater flexibility and anatomical differences typically seen in women.

In contrast, male players demonstrated superior performance in the Trunk Stability Push-Up and Rotary Stability tests, likely due to more efficient activation and coordination of key stabilizing muscle groups, including the hip adductors, rotators, abdominal muscles, spinal stabilizers, and the quadratus lumborum (35-37).

Johnson et al. (2023) investigated sex-related differences in FMS scores among Reserve Officers' Training Corps (ROTC) cadets. The findings revealed that females outperformed males on both the Inline Lunge and Active Straight Leg Raise tests. Conversely, males demonstrated significantly higher scores on the Trunk Stability Push-Up test. In a similar vein, these results align with our study as well. These differences may be attributed to increased mobility in the ankle and hamstrings, as well as reduced upper-body muscle mass in females compared to males. Consequently, these sex-related differences should be taken into account when designing tailored exercise programs for males and females (38).

In the present study, no significant differences were observed between females and males in the total FMS score. The literature on FMS presents conflicting evidence regarding sex differences in total scores; some studies (39-41), have reported that males demonstrate higher quality movement patterns than females. However, more recent research suggests that females tend to perform better when composite FMS scores are evaluated (42) . Although there may be general differences in physical performance between males and females due to biological factors such as muscle mass and hormonal variations (43), it is important to recognize that individuals of each sex can exhibit a wide range of abilities and capabilities (41).

Additionally, several studies have compared FMS scores between male and female athletes across different populations. According to Hamil et al. (2021), female soccer players exhibited

higher FMS scores and greater injury rates than male players. They suggest that this discrepancy may justify increasing the FMS cutoff score for females as a predictor of injury. However, further research is needed due to the small sample size of their study (٤٤). According to Ransdell and Murray (2016), the higher incidence of injuries among females can be attributed to a range of contributing factors (٤٦ ,٤٥). These include reduced muscle fiber size and cross-sectional area (47), a narrower skeletal structure (48), decreased activation of the gluteus medius muscle (35), as well as the vastus medialis oblique and vastus lateralis (49), diminished neuromuscular coordination (50, 51), and less core stability (37, 50). In contrast, Amjad et al. (2023) found that female footballers scored lower on the FMS tests than male footballers. They concluded that there was a significant association between FMS composite scores and injury occurrence, suggesting that the FMS results for both genders may be linked to the likelihood of injury (14).

The results of the current study revealed significant differences in lower extremity performance between male and female soccer players, specifically in the Single Hop for Distance (P = 0.001) and Cross-over Hop for Distance (P = 0.001). Analysis of the mean scores indicated that male players outperformed female players in both tests. These findings suggest that male athletes tend to exhibit superior explosive power and strength in the lower body, likely due to greater muscle mass and differences in hormonal profiles, which are known to contribute to enhanced performance in power-related tasks such as hopping and distance-based tests (37, 52). In contrast, when comparing upper extremity performance, significant differences were also observed between male and female soccer players. Specifically, male athletes performed better in the Davies test (P = 0.001) and upper extremity dynamic balance in the superolateral direction (P = 0.026). These results align with previous studies that indicate males generally demonstrate greater upper body strength and neuromuscular control, which is often attributed to increased muscle mass in the upper body and higher testosterone levels (37, 52).

Our findings show that males performed superior in all physical ability tests. Reference data from youth to adulthood typically indicate that males possess greater musculoskeletal strength and power than females, which aligns with the observed results in the physical performance tests. In support of this, O'Brien-Smith et al. (20١٩), the performance of male and female football players aged 9 to 18 was compared. It was reported that male players outperformed females in factors such as individual skills, vertical jump, sprinting, agility, and dribbling power. The study attributed these differences to gender-based anthropometric characteristics, which were considered among the reasons for the superiority of male players (18). Similarly, Schons et al. (2023) compared the anthropometric profiles and physical performance of professional male and female soccer players. It confirmed that men generally perform better than women in physical tests (53).

In another study by Arundale et al. (2020), it was found that female athletes are at greater risk for lower extremity injuries, including Anterior Cruciate Ligament (ACL) injuries. Diagnostic test scores for lower limb movement pattern deficits such as increased knee valgus and decreased knee flexion were more pronounced in females compared to males. These findings were attributed to lower limb muscular weakness in female athletes relative to their male counterparts (54). Consequently, it could help justify the findings of specific studies that suggest women exhibit weaker performance compared to men.

Based on the findings of the present study, it is recommended that coaches, movement specialists, and sports medicine teams take gender differences in physical performance into account when designing training programs and injury prevention strategies.

Focusing on core stability exercises in female athletes due to a higher prevalence of trunk instability and incorporating flexibility-enhancing exercises in male athletes to reduce muscular tightness and improve range of motion can play a significant role in reducing injury risk and enhancing athletic performance. For example, exercises such as the Plank with leg lift can help women improve core stability and reduce lower limb stress, while the World's Greatest Stretch can aid men in increasing hip joint and hamstring flexibility, thereby improving mobility and functional performance.

Furthermore, future research should carefully control for potential confounding variables such as training history, injury history, and anthropometric characteristics. For example, athletes with longer and more consistent training experience typically demonstrate greater strength, endurance, and movement efficiency, factors that can influence their test performance and susceptibility to injury, to elucidate better the role of functional assessment tools like the FMS in predicting injury susceptibility and improving movement efficiency.

5. Conclusions

This study demonstrates the necessity of gender-specific training programs, as they are crucial in enhancing performance and reducing the risk of injury. According to the findings, it is recommended that flexibility and mobility exercises be implemented for male soccer players and core stability and strength training for female soccer players. These suggestions stem from the observed FMS tests. Additionally, male athletes outperformed females in performance tests, likely due to biomechanical and neuromuscular variations between the sexes. In this regard, future research should investigate the underlying mechanisms contributing to these differences, considering individual factors such as playing experience and fitness level.

6. Limitations of the Study

Despite the researchers' efforts to control various aspects of the study, several limitations remain. One major limitation is the small sample size, which reduces the generalizability of the findings. Increasing the number of participants in future research is essential to enhance statistical power and obtain more reliable results. Another limitation is that data collection occurred only once during the season, potentially overlooking variations in movement quality and performance across different training periods. Therefore, Future studies should assess FMS and physical performance before and after training phases to clarify how changes in movement patterns affect performance outcomes over time.

Declaration of Interest Statement

Ethics approval

This study was approved by the Sport Sciences Research Institute of Iran. Approval code**:** SSRI.REC-2310-2493
Date of approval: 23thOcrober 2023.

Consent for publication

All authors have given their consent for the publication of this manuscript.

Availability of data and materials

All relevant data and materials are included within the article. Additional data can be made available upon reasonable request.

Credit authorship contribution statement

AZ: Data curation, writing the original draft. **HP**: Conceptualization, Supervision, and paper editing. **RH**: writing – review & editing. **AS**: review and editing. **AQ**: Review and editing.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Fundings

This research did not receive any specific funding.

Acknowledgments

The authors sincerely thank everyone who supported them throughout the study.

References

.١ Griffin J, Larsen B, Horan S, Keogh J, Dodd K, Andreatta M, et al. Women's Football: An Examination of Factors That Influence Movement Patterns. J Strength Cond Res. 2020;34(8):2384–93.

.٢ Marqués Jiménez D, Calleja González J, Arratibel Imaz I, Delextrat A, Terrados Cepeda N. Fatigue and recovery in soccer: Evidence and challenge. Open Sports Sciences Journal. 2017.

.٣ Samudra AD, Purwanto B, Utomo DN. Differences in Limb Muscle Strength Affecting Vertical Jump Heights in Soccer Players after Chronic Ankle Injury. Journal of Modern Rehabilitation. 2024.

.٤ Polinder S, Haagsma J, Panneman M, Scholten A, Brugmans M, Van Beeck E. The economic burden of injury: Health care and productivity costs of injuries in the Netherlands. Accid Anal Prev. 2016;93:92.١٠٠–

.٥ Asgari M, Alizadeh S, Sendt A, Jaitner T. Evaluation of the Functional Movement Screen (FMS) in Identifying Active Females Who are Prone to Injury. A Systematic Review. Sports Med Open. 2021;7(1):85.

.٦ Kramer TA, Sacko RS, Pfeifer CE, Gatens DR ,Goins JM, Stodden DF. The association between the functional movement screen™, y-balance test, and physical performance tests in male and female high school athletes. Int J Sports Phys Ther. 2019;14(6):911–9.

.٧ Lim KH, Seo TB, Kim YP. Relationship between movement dysfunctions and sports injuries according to gender of youth soccer player. J Exerc Rehabil. 2020;16(5):427–31.

.٨ Pardos-Mainer E, Lozano D, Torrontegui-Duarte M, Cartón-Llorente A, Roso-Moliner A. Effects of Strength vs. Plyometric Training Programs on Vertical Jumping, Linear Sprint and Change of Direction Speed Performance in Female Soccer Players: A Systematic Review and Meta-Analysis. Int J Environ Res Public Health. 2021;18.(٢)

.٩ Cook G, Burton L, Kiesel K, Bryant MF, Rose G. Movement :Functional Movement Systems : Screening, Assessment, and Corrective Strategies: On Target Publications; 2010.

.١٠ Cook G, Burton L, Hoogenboom B. Pre-participation screening: the use of fundamental movements as an assessment of function - part 1. N Am J Sports Phys Ther. 2006;1(2):62–72.

.۱۱ Farley JB, Barrett LM, Keogh JWL, Woods CT, Milne N. The relationship between physical fitness attributes and sports injury in female, team ball sport players: a systematic review. Sports Med Open. 2020;6(1):45.

.۱۲ Mahrokh Moghadam A, Zarei M, Mohammadi F. Effect of an eight-week core stability training program on the functional movement screen test scores in elite goalball players. Physical Treatments-Specific Physical Therapy Journal. 2021;11(1):55–62.

.۱۳ Martín-Moya R, Rodríguez-García L, Moreno-Vecino B, Clemente FM, Liñán González A, González-Fernández FT. Differences and relationship in functional movement screen (FMS™) scores and physical fitness in males and female semi-professional soccer players. PeerJ. 2:۱۱;۰۲۳e16649.

.۱۴ Amjad F, Latif D, Hanif S, Bashir K, Batool R, Khalid U. Functional Movement Screen Differences in Male and Female Footballers and as an Injury Prevention Tool: Functional Movement Screen Differences. Pakistan Journal of Health Sciences.۴–۲۰۲۳:۳۰ .

.۱۵ Emmonds S, Morris R, Murray E, Robinson C, Turner L, Jones B. The influence of age and maturity status on the maximum and explosive strength characteristics of elite youth female soccer players. Science and Medicine in Football. 2017;1.۱۵–۲۰۹:(۳)

.۱۶ Manson SA, Brughelli M, Harris NK. Physiological characteristics of international female soccer players. J Strength Cond Res. 2014;28(2):308–18.

.۱۷ Mascherini G, Levi Micheli M, Serafini S, Politi C, Bianchi E, Cebrián-Ponce Á, et al. Raw bioelectrical data and physical performance in track and field athletes: Are there differences between the sexes in the relationship? Heliyon. 2024;10(15):e35754.

.۱۸ O'Brien-Smith J, Bennett K, Fransen J, Smith M. Same or different? A comparison of anthropometry, physical fitness and perceptual motor characteristics in male and female youth soccer players. Science and Medicine in Football. 2019;4(6):1–8.

.۱۹ Datson N, Weston M, Drust B, Gregson W, Lolli L. High-intensity endurance capacity assessment as a tool for talent identification in elite youth female soccer. J Sports Sci. 2020;38(11-12):1313–9.

.۲۰ Cook G, Burton L, Hoogenboom BJ, Voight M. Functional movement screening: the use of fundamental movements as an assessment of function-part 2. Int J Sports Phys Ther. 2014;9(4):549–63.

.۲۱ Bonazza NA, Smuin D, Onks CA, Silvis ML, Dhawan A. Reliability, Validity, and Injury Predictive Value of the Functional Movement Screen: A Systematic Review and Meta-analysis. Am J Sports Med. 2017;45(3):725–32.

.۲۲ Karimizadeh Ardakani M, Soroush Fard Z, Amirizadeh F, Naderifar H. Effect of thoracic hyper-kyphosis posture on upper extremity function of female students. Journal of Rehabilitation Sciences & Research. 2022;9(1):30–5.

.۲۳ Narducci E, Waltz A, Gorski K, Leppla L, Donaldson M. The clinical utility of functional performance tests within one-year post-acl reconstruction: a systematic review. Int J Sports Phys Ther. 2011;6(4):333–42.

.۲۴ Noyes FR, Barber SD, Mangine RE. Abnormal lower limb symmetry determined by function hop tests after anterior cruciate ligament rupture. The American journal of sports medicine. 1991;19(5):513–8.

.۲۵ Ross MD, Langford B, Whelan PJ. Test-retest reliability of 4 single-leg horizontal hop tests. ournal of strength and conditioning research. 2002;16(4):617–22.

۲۶. Reid A, Birmingham TB, Stratford PW, Alcock GK, Giffin JR. Hop testing provides a reliable and valid outcome measure during rehabilitation after anterior cruciate ligament reconstruction. Phys Ther. 2007;87(3):337–49.

۲۷. Noyes FR, Barber SD, Mangine RE. Abnormal lower limb symmetry determined by function hop tests after anterior cruciate ligament rupture. Am J Sports Med. 1991;19(5):513–8.

۲۸. Ross MD, Langford B, Whelan PJ. Test-retest reliability of 4 single-leg horizontal hop tests. J Strength Cond Res. 2002;16(4):617–22.

۲۹. Gorman PP, Butler RJ, Plisky PJ, Kiesel KB. Upper Quarter Y Balance Test: reliability and performance comparison between genders in active adults. J Strength Cond Res. 2012;26(11):3043–8.

۳۰. Hazar Z, Ulug N, Yuksel I. Upper quarter y-balance test score of patients with shoulder impingement syndrome. Orthopaedic journal of sports medicine. 2014;2(11_suppl3):2325967114S00275.

۳۱. Marques VB, Medeiros TM, de Souza Stigger F, Nakamura FY, Baroni BM .The Functional Movement Screen (FMS™) in Elite Young Soccer Players Between 14 and 20 Years: Composite Score, Individual-Test Scores, and Asymmetries. Int J Sports Phys Ther. 2017;12(6):977–85.

۳۲. Medeiros DM, Miranda LLP, Marques VB, de Araujo Ribeiro-Alvares JB, Baroni BM. Accuracy of the Functional Movement Screen (FMS™) active straight leg raise test to evaluate hamstring flexibility in soccer players. Int J Sports Phys Ther. 2019;14(6):877–84.

۳۳. Minick KI, Kiesel KB, Burton L, Taylor A, Plisky P ,Butler RJ. Interrater reliability of the functional movement screen. J Strength Cond Res. 2010;24(2):479–86.

۳۴. Shultz R, Anderson SC, Matheson GO, Marcello B, Besier T. Test-retest and interrater reliability of the functional movement screen. J Athl Train. 2013;48(3):331–6.

۳۵. Hart JM, Garrison JC, Kerrigan DC, Palmieri-Smith R, Ingersoll CD. Gender differences in gluteus medius muscle activity exist in soccer players performing a forward jump. Res Sports Med. 2007;15(2):147–55.

۳۶. Kernozek TW, Torry MR, Iwasaki M. Gender differences in lower extremity landing mechanics caused by neuromuscular fatigue. Am J Sports Med. 2008;36(3):554–65.

۳۷. Zazulak BT, Hewett TE, Reeves NP, Goldberg B, Cholewicki J. Deficits in neuromuscular control of the trunk predict knee injury risk: a prospective biomechanical-epidemiologic study. Am J Sports Med. 2007;35(7):1123–30.

۳۸. Johnson QR, Scraper J, Lockie R, Orr RM, Dawes JJ. Sex-related Differences in Functional Movement Screen Scores Among Reserve Officers' Training Corps Cadets. Mil Med. 2023;188(1-2):e152–e7.

۳۹. Chimera NJ, Smith CA, Warren M. Injury history, sex, and performance on the functional movement screen and Y balance test. J Athl Train. 2015;50(5):475–85.

۴۰. Paszkewicz JR, McCarty CW, Van Lunen BL .Comparison of functional and static evaluation tools among adolescent athletes. J Strength Cond Res. 2013;27(10):2842–50.

۴۱. Domaradzki J, Koźlenia D. Reliability of Functional Movement Screen and sexual differentiation in FMS scores and the cut-off point among amateur athletes. 2020.

۴۲. Thomas K, Holmes L, Wolf D, editors. Gender differences in functional movement screening scores in men’s and women’s collegiate tennis. International Journal of Exercise Science: Conference Proceedings; 2022.

۴۳. Ansdell P, Thomas K, Hicks KM, Hunter SK, Howatson G, Goodall S. Physiological sex differences affect the integrative response to exercise: acute and chronic implications. Exp Physiol. 2020;105(12):2007–21.

۴۴. Hamil JA, Howell SM, Deldin AR. Functional Movement Screen differences between male and female collegiate soccer players. Journal of Physical Education. 2021;8(2):34–9.

۴۵. Ransdell L, Murray T. Functional Movement Screening. Strength and Conditioning Journal. 2016;38(2):40–8.

۴۶. Ransdell LB, Wells CL .Sex Differences in Athletic Performance. Women in Sport and Physical Activity Journal. 1999;8(1):55–81.

۴۷. Sale DG, MacDougall JD, Alway SE, Sutton JR. Voluntary strength and muscle characteristics in untrained men and women and male bodybuilders. J Appl Physiol (1985). 1987;62(5):1786–93.

۴۸. Strength N-N, Association C. Essentials of strength training and conditioning: Human kinetics; 2021.

۴۹. Kim M-H, Yoo W-G, Yi C-H. Gender differences in the activity and ratio of vastus medialis oblique and vastus lateralis muscles during drop landing. Journal of Physical Therapy Science. 2009;21(4):325–9.

۵۰. Brophy RH, Chiaia TA, Maschi R, Dodson CC, Oh LS, Lyman S, et al. The core and hip in soccer athletes compared by gender. Int J Sports Med. 2009;30(9):663–7.

۵۱. Hughes G, Watkins J, Owen N. Gender differences in lower limb frontal plane kinematics during landing. Sports Biomech. 2008;7(3):333–41.

۵۲. Storer TW, Basaria S, Traustadottir T, Harman SM, Pencina K, Li Z, et al. Effects of Testosterone Supplementation for 3 Years on Muscle Performance and Physical Function in Older Men. J Clin Endocrinol Metab. 2017;102(2):583–93.

۵۳. Schons P, Preissler A, Oliveira R, Brito J, Clemente F, de Vargas G, et al. Comparisons and correlations between the anthropometric profile and physical performance of professional female and male soccer players: Individualities that should be considered in training. International Journal of Sports Science & Coaching. 2023;18(6):2004–14.

۵۴. Arundale AJH, Kvist J, Hägglund M, Fältström A. Jump performance in male and female football players. Knee surgery, sports traumatology, arthroscopy : official journal of the ESSKA. 2020;28(2):606–13.